

RESEARCH ARTICLE

On the construction of the KP line-solitons and their interactions

Sarbarish Chakravarty^{a,1}, Tim Lewkow^a and Ken-ichi Maruno^b^aDepartment of Mathematics, University of Colorado, Colorado Springs, CO 80933^bDepartment of Mathematics, University of Texas-Pan American, Edinburg, TX 78539

(Received 00 Month 200x; in final form 00 Month 200x)

The line-soliton solutions of the Kadomtsev–Petviashvili (KP) equation are investigated in this article using the τ -function formalism. In particular, the Wronskian and the Grammian forms of the τ -function are discussed, and the equivalence of these two forms are established. Furthermore, the interaction properties of two special types of 2-soliton solutions of the KP equation are studied in details.

Keywords: KP equation; line-solitons; τ -function

AMS Subject Classifications: 37K10, 37K35, 37K40

1. Introduction

The eponymous Kadomtsev–Petviashvili (KP) equation

$$(-4u_t + u_{xxx} + 6uu_x)_x + 3\beta u_{yy} = 0, \quad (1)$$

discovered in 1970, describes the dynamics of small-amplitude, long wavelength, solitary waves in two dimensions (xy -plane) [1], and arises in the study of water waves, plasma and various other areas of physical significance (see e.g., [2] for a review). In equation (1), $u = u(x, y, t)$ is the wave amplitude, the subscripts denote partial derivatives with respect to x, y, t , and $\beta = \pm 1$. Throughout this article, equation (1) with $\beta = 1$, which corresponds to the negative-dispersion KP equation (KP II) will be considered, and will be referred to as the KP equation.

The KP equation is a completely integrable system whose underlying mathematical properties have been extensively studied during the last four decades. They are well documented in several monographs including (but not limited to) [3, 4, 5, 6]. These properties include the existence of multi-soliton and periodic solutions and the Lax representation of the inverse scattering transform. A major breakthrough in the KP theory occurred in 1981 when Sato [7] formulated the KP equation in terms of an infinite dimensional Grassmann manifold known as the Sato universal Grassmannian. A finite dimensional version of the Sato theory corresponding to the real Grassmannian $\text{Gr}(N, M)$ (the set of N -dimensional subspaces of $\mathbb{R}^M$) leads to a simple algebraic construction of a special class of solitary wave solutions of the KP equation called the *line-soliton* solutions. These are real, non-singular solutions which decay exponentially in the xy -plane except along certain directions. Specifically, such a solution is localized along two distinct sets of rays (referred to

¹Corresponding author. Email: chuck@math.uccs.edu

as line solitons) as $y \rightarrow \pm\infty$, and form spatial interaction patterns in the finite region of the xy -plane.

The simplest example of a line-soliton is the one-soliton solution of KP given by

$$u(x, y, t) = \alpha \operatorname{sech}^2(\mathbf{k} \cdot \mathbf{r} + \omega t + \theta_0), \quad (2)$$

which is a traveling wave with $\mathbf{r} := (x, y)$, amplitude $\alpha = \frac{1}{2}(k_2 - k_1)^2$, wave vector $\mathbf{k} := (k_x, k_y) = \frac{1}{2}(k_2 - k_1, k_2^2 - k_1^2)$ and frequency $\omega = \frac{1}{2}(k_2^3 - k_1^3)$, where k_1, k_2 are distinct real parameters with $k_1 < k_2$. Clearly, the soliton amplitude depends on the wave vector $\mathbf{k}$, which together with the frequency ω satisfy the soliton dispersion relation

$$4\omega k_x = 4k_x^4 + 3k_y^2.$$

The solitary wave-form given by (2) is localized in the xy -plane along a line which makes an angle ψ measured counterclockwise from the y -axis where

$$\tan \psi = k_y/k_x = k_1 + k_2, \quad -\frac{\pi}{2} < \psi < \frac{\pi}{2}.$$

A one-soliton solution is shown in Figure 1(a). Since the one-soliton solution is characterized by two real parameters k_1, k_2 , it is convenient to denote this solution simply as the $[1, 2]$ -soliton. Note that when $k_1 + k_2 = 0$, the solution in (2) becomes y -independent and reduces to the one-soliton solution of the Korteweg-de Vries (KdV) equation.

The soliton interactions of the KP equation were originally described via a 2-soliton solution with a “X”-shape pattern in the plane formed by the intersection of two line solitons but with a parallel shift of the two lines at the intersection (the phase shift). This 2-soliton solution (see Figure 1(c)) referred to as the O-type soliton (“O” stands for *original*) and its N -soliton generalization were obtained independently by several authors using integral equations [8] and other algebraic methods [9, 25]. In 1977, Miles [11] pointed out that the O-type 2-soliton solution becomes singular if the angle of the intersection is smaller than a certain critical value. As the angle approaches the critical value from above, the 2-soliton phase shift tends to infinity, and at the critical angle the O-type solution degenerates to a “Y-shape” with only three line solitons interacting resonantly (see also [12]). It turns out that such Y-shape interacting wave-forms are also exact solutions of the KP equation [13, 14]. Apart from the ones mentioned thus far, no other soliton solutions of the KP equation were known for quite some time until recently when more general types of resonant and non-resonant line-soliton solutions were reported in several works including [15, 16, 17, 18]. During the last 5 years, considerable progress has been made toward the problem of classifying all exact line-soliton solutions of the KP equation [19, 20, 21, 22]. These studies have revealed a large variety of soliton solutions which were totally overlooked in the past. Generically, these solutions of KP consist of two distinct sets of line solitons of different amplitudes and along different directions in the xy -plane as $y \rightarrow \pm\infty$. Locally, each line soliton denoted as the $[i, j]$ -soliton has the form of a one-soliton solution as in (2), parametrized by two distinct real parameters $k_i < k_j, i < j$. The soliton amplitude is given by $\alpha_{ij} = \frac{1}{2}(k_j - k_i)^2$ and the soliton angle ψ_{ij} (counterclockwise from the y -axis) satisfies $\tan \psi_{ij} = k_i + k_j$.

There are several direct and algebraic methods to construct the KP line-soliton solutions which are usually derived from either a Wronskian or from a Gram determinant (Gramian). In principle any given soliton solution can be represented in either form, although this fact has been explicitly shown for only the one-soliton and the O-type soliton solutions. For more general line-soliton solutions, it is often more difficult to impose regularity conditions on the solution when it is represented in one particular form than the other, thus rendering one solution generating method less efficient than another. In this note, we dis-




Figure 1. Line-soliton solutions of KP: (a) a one-soliton solution with $(k_1, k_2) = (0.5, 1)$ at $t = 0$, ψ is the soliton angle; (b) a Miles's Y-shape soliton solution with $(k_1, k_2, k_3) = (-1, 0, \frac{1}{2})$ at $t = 0$; (c) an O-type 2-soliton with $(k_1, \dots, k_4) = (-1, -\frac{1}{2}, \frac{1}{2}, 2)$ at $t = 0$. In each figure the dominant exponentials in the τ -function (see section 2) in different regions of the xy -plane are labeled by the indices in parentheses, and the asymptotic line solitons are labeled by the index pair in square brackets.

cuss both the Wronskian and the Grammian forms of the general line-soliton solutions, and establish the equivalence between the two representations in an explicit fashion. That is, we show how to derive one form of the solution from the other, and vice-versa. Another purpose of this paper is to describe the interaction properties of the various types of line-soliton solutions. In particular, we present a detailed discussion of two distinct types of two-soliton solutions both of which form an “X”-shape pattern on the xy -plane but interact in a significantly different way.

The paper is planned as follows. In Section 2, we introduce the τ -function of the KP equation and present its Wronskian and Grammian forms generating the line-soliton solutions. Then the equivalence of these two representations of the τ -function is established. Section 3 is devoted to a brief description of the distinct types of the 2-soliton solutions, followed by a detailed discussion of the nonlinear interaction properties of the O-type and P-type 2-soliton solutions. We conclude the paper with a brief summary and possible significance of the results.

2. The KP τ -function

The most convenient representation for the line-soliton solutions is via the τ -function which plays an important role in the mathematical theory of the KP equation [6, 7]. The solution $u(x, y, t)$ of (1) can be expressed as

$$u(x, y, t) = 2(\ln \tau)_{xx}, \quad (3)$$

where $\tau(x, y, t)$ is the KP τ -function which is defined up to an exponential factor that is linear in x, y and t .

2.1. The Wronskian form of the τ -function

The τ -function can be expressed as the Wronskian determinant [7, 24, 25]

$$\tau(x, y, t) = \text{Wr}(f_1, \dots, f_N) = \begin{pmatrix} f_1 & f_2 & \cdots & f_N \\ f_1^{(1)} & f_2^{(1)} & \cdots & f_N^{(1)} \\ \vdots & \vdots & \ddots & \vdots \\ f_N^{(N-1)} & f_2^{(N-1)} & \cdots & f_N^{(N-1)} \end{pmatrix}, \quad (4)$$

where $f^{(i)}$ denotes the i^{th} partial derivative with respect to x , and where the functions $\{f_n\}_{n=1}^N$ form a set of linearly independent solutions of the linear system

$$f_y = f_{xx}, \quad f_t = f_{xxx}. \quad (5)$$

A remarkable fact is that the KP equation simply turns into a determinant identity if one substitutes (3) into (1) and uses the Wronskian form of $\tau(x, y, t)$ given by (4) (see e.g., [6, 23, 25] for details). This implies that *any* linearly independent set of solutions of the linear system (5) will give rise to a solution of the KP equation. Hence it is possible to generate a large class of solutions in this way. In particular, the line-soliton solutions are obtained from the choice

$$f_n(x, y, t) = \sum_{m=1}^M a_{nm} e^{\theta_m}, \quad n = 1, 2, \dots, N, \quad (6)$$

where $\theta_m(x, y, t) = k_m x + k_m^2 y + k_m^3 t + \theta_{0m}$ with distinct real parameters: $k_1 < k_2 < \dots < k_M$ and real constants $\{\theta_{0m}\}_{m=1}^M$. The coefficients $(a_{nm}) =: A$ define an $N \times M$ constant matrix which is of rank N due to the linear independence of the functions $\{f_n\}_{n=1}^N$. Then (4) can be expressed as

$$\tau(x, y, t) = \sum_{1 \leq m_1 < \dots < m_N \leq M} A(m_1, \dots, m_N) \exp[\theta(m_1, \dots, m_N)] \prod_{1 \leq s < r \leq N} (k_{m_r} - k_{m_s}), \quad (7)$$

by expanding the Wronskian determinant using Binet-Cauchy formula. In above, $\theta(m_1, \dots, m_N) := \theta_{m_1} + \theta_{m_2} + \dots + \theta_{m_N}$, and $A(m_1, \dots, m_N)$ is the maximal minor, i.e., the determinant of the $N \times N$ sub-matrix of A obtained from columns $1 \leq m_1 < \dots < m_N \leq M$. Note that the N linearly independent rows of the coefficient matrix A span an N -dimensional subspace W of $\mathbb{R}^M$ so that A can be regarded as a point of the real Grassmannian $\text{Gr}(N, M)$. A different choice of basis for W amounts to performing row operations: $A \rightarrow GA$, $G \in \text{GL}(N, \mathbb{R})$, which changes the τ -function in (7) simply by a scale factor: $\tau \rightarrow \det(G)\tau$ but leaves the KP solution $u = 2(\ln \tau)_{xx}$ invariant. Consequently the coefficient matrix A can be canonically chosen in the reduced row echelon form (RREF) which has a distinguished set of pivot columns such that the restriction of A to this set corresponds to the $N \times N$ identity matrix. Given any $N \times M$ matrix A of rank N , its unique RREF gives a standard coordinate for the corresponding point in $\text{Gr}(N, M)$. It is in this sense that the space of all solutions of KP generated by the τ -function given by (7) can be identified with the real Grassmann manifold $\text{Gr}(N, M)$. In general, such solutions are not all line-solitons as they can be singular where the τ -function becomes zero. The regularity of the solutions in the entire xy -plane and for all values of t is achieved by imposing a further restriction on the coefficient matrix A , namely that all of its $N \times N$ maximal minors be non-negative. Such matrices are called *totally non-negative* matrices, they represent the totally non-negative Grassmannian $\text{Gr}^+(N, M)$ which forms a closed subset in $\text{Gr}(N, M)$, and which identifies the space of all line-soliton solutions of the KP equation.

Some simple examples of τ -functions corresponding to some of the solutions mentioned in Section 1 are listed below.

One-soliton solution: In this case, one chooses a single function in (6) of the form $f_1 = e^{\theta_1} + e^{\theta_2}$. The coefficient matrix is simply $A = \begin{pmatrix} 1 & 1 \end{pmatrix}$, and $\tau = f_1$. Then equation (3) yields the one-soliton solution given by (2) in Section 1 with $\theta_0 = \theta_{02} - \theta_{01}$. Note that since $k_1 < k_2$, $\tau \sim e^{\theta_1} \gg e^{\theta_2}$ and $u(x, y, t) \sim 0$ as $x \rightarrow -\infty$. A similar argument shows that u is also exponentially small as $x \rightarrow \infty$. In this case, one finds from the exact expression in (2) that the solution is localized in the xy -plane along the line $\theta_1 = \theta_2$ (for

fixed t) where the two exponentials in the τ -function are precisely in balance (see Figure 1(a)). It turns out that even for general line-solitons, the solution $u(x, y, t)$ is localized along certain lines in the xy -plane where exactly two exponential terms in the τ -function of equation (7) are in balance, and they dominate over all other exponential terms. This principle of *dominant balance* was implemented in Refs [17, 19, 20, 21] to identify the asymptotic line solitons $[i, j]$ associated with each line-soliton solution of the KP equation.

Miles Y-shape solution: The τ -function for this solution is a sum of three exponential functions, i.e., $\tau = f_1 = e^{\theta_1} + e^{\theta_2} + e^{\theta_3}$ with the coefficient matrix $A = \begin{pmatrix} 1 & 1 & 1 \end{pmatrix}$. Applying the dominant balance principle mentioned above, it is possible here to determine the dominant exponentials and analyze the structure of the solution in the xy -plane. One finds that the solution $u(x, y, t)$ is localized along three lines corresponding to the line soliton $[1, 3]$ for $y \gg 1$, and line solitons $[1, 2]$, $[2, 3]$ for $y \ll 1$ as illustrated in Figure 1(b). For $i, j \in \{1, 2, 3\}$, each line soliton $[i, j]$, $i < j$ is locally of the form of the one-soliton solution (2) with distinct parameters k_i, k_j . As mentioned in Section 1, the Miles solution represents a resonant solution of three line solitons. The resonant condition among those three line-solitons is given by

$$\mathbf{k}_{13} = \mathbf{k}_{12} + \mathbf{k}_{23}, \quad \omega_{13} = \omega_{12} + \omega_{23},$$

where $\mathbf{k}_{ij} = \frac{1}{2}(k_j - k_i, k_j^2 - k_i^2)$ and $\omega_{ij} = \frac{1}{2}(k_j^3 - k_i^3)$ are respectively, the wave vector and frequency of the line soliton $[i, j]$, satisfying the soliton dispersion relation given in Section 1.

O-type 2-soliton solution: Here the τ -function is a Wronskian of two linearly independent solutions of (5). These solutions are obtained from (6) by choosing $N = 2$, $M = 4$, and the coefficient matrix

$$A = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix},$$

such that $f_1 = e^{\theta_1} + e^{\theta_2}$ and $f_2 = e^{\theta_3} + e^{\theta_4}$. Note that A is a totally non-negative matrix whose 2×2 maximal minors are: $A(1, 2) = A(3, 4) = 0$, and $A(1, 3) = A(1, 4) = A(2, 3) = A(2, 4) = 1$. From (7), the resulting τ -function is given by

$$\tau = (k_3 - k_1)e^{\theta(1,3)} + (k_4 - k_1)e^{\theta(1,4)} + (k_3 - k_2)e^{\theta(2,3)} + (k_4 - k_2)e^{\theta(2,4)}, \quad \theta(i, j) := \theta_i + \theta_j.$$

Figure 1(c) illustrates the regions in the xy -plane where each of the four exponential terms in the above τ -function are dominant. The line solitons corresponding to the solution $u = 2(\ln \tau)_{xx}$ are $[1, 2]$ and $[3, 4]$ for $|y| \rightarrow \infty$; these are localized along the directions where a pair of dominant exponential terms are in balance.

2.2. The Grammian form of the τ -function

The Grammian construction of the line-soliton solutions will be described next. This form of the solution arises from the so called binary Darboux transformation of the KP equation (see e.g., [24]). In this case, the τ -function can be expressed as a determinant of an $N \times N$ matrix as follows:

$$\tau = \det(\Omega), \quad \Omega_{ij} = \int_{\Gamma} f_i g_j dx - (f_i g_{jx} - f_{ix} g_j) dy, \quad i, j = 1, 2, \dots, N, \quad (8)$$

where $\{f_i\}_{i=1}^N$ solve (5) and $\{g_j\}_{j=1}^N$ solve the (formal) adjoint system of (5), namely,

$$-g_y = g_{xx}, \quad g_t = g_{xxx}.$$

It is easily verified from (5) and its adjoint that $d\Omega_{ij}$ is an exact differential, hence the line integral above can be evaluated along any suitable curve Γ in the xy -plane such that the integral converges. Like the Wronskian form, it can be shown that the τ -function in (8) yields a solution of the KP equation via (3) for any choice of the sets of functions $\{f_i\}_{i=1}^N$ and $\{g_j\}_{j=1}^N$ [6]. For the line-soliton solutions, these are chosen as linear combinations of exponentials (cf. (6)),

$$f_i = \sum_{n=1}^r b_{in} e^{\phi(p_n)}, \quad g_j = \sum_{n=1}^N \hat{b}_{jn} e^{-\phi(q_n)}, \quad i, j = 1, \dots, N, \quad (9)$$

where $\phi(k) := kx + k^2y + k^3t + \phi_0(k)$, and all the parameters $p_1, \dots, p_r, q_1, \dots, q_N$ are distinct and real. Note that $N \neq r$, in general. Thus, $B := (b_{mn})$ is an $N \times r$ matrix whereas $\hat{B} := (\hat{b}_{mn})$ is an invertible, $N \times N$ matrix. Substituting (9) into the matrix elements Ω_{ij} in (8), and evaluating the line integral along a path Γ from $(-\sigma\infty, y)$ to (x, y) parallel to the x -axis, yields

$$\Omega_{ij}(x, y, t) = c_{ij} + \int_{\Gamma} f_i g_j dx = c_{ij} + \sum_{m=1}^r \sum_{n=1}^N b_{im} \hat{b}_{jn} \frac{e^{\phi(p_m) - \phi(q_n)}}{p_m - q_n}, \quad (10)$$

where $\sigma = \text{sgn}(p_m - q_n) = \pm 1$ and c_{ij} are arbitrary constants. Choosing the constants $c_{ij} = \delta_{ij}$, where δ_{ij} is the Kröner symbol ($\delta_{ij} = 0, i \neq j, \delta_{ii} = 1$), ensures that the matrix Ω is of rank N so that $\tau \neq 0$. Using (10), the τ -function in (8) can be expressed in the form

$$\tau = \det(I + BF\hat{B}^T) = \det(I + CF), \quad (11)$$

where $C = \hat{B}^T B$ is an $N \times r$ matrix of constant coefficients and F is an $r \times N$ matrix whose entries are given by

$$F_{mn} = \frac{e^{\phi(p_m) - \phi(q_n)}}{p_m - q_n}, \quad m = 1, 2, \dots, r, \quad n = 1, 2, \dots, N.$$

The Grammian form given by the first equality in (11) is not unique as it is possible to obtain the same τ -function from a different choice of the matrices B and $\hat{B}$. In particular, $\hat{B}$ need not be a square matrix although there is always a canonical choice of an $N \times N$ matrix $\hat{B}$ that leads to the second equality in (11). We will not discuss the details of the gauge freedom underlying the choice of B and $\hat{B}$ in this article.

Equation (11) is the canonical Grammian form for the line-soliton solutions, special cases of which arise from several direct methods of constructing solutions including the Hirota method [6], direct linearization [4] and dressing techniques [8]. For example, the O-type N -soliton solution is obtained by setting $r = N$ and $C = I$, the latter being the $N \times N$ identity matrix. In this case the resulting τ -function $\tau = \det(I + F)$ is positive for all x, y, t if the parameters are ordered as $p_N < p_{N-1} < \dots < p_1 < q_1 < q_2 < \dots < q_N$. Then the corresponding line-soliton solution is non-singular. More general line-soliton solutions can also be generated from the Grammian form (11) by making appropriate choices for the matrix C . However, in comparison with the Wronskian form, here it is less clear how to impose regularity conditions on the obtained solution. Recall that in

the Wronskian construction the ordering $k_1 < k_2 < \dots < k_M$ of the parameters and the totally non-negative coefficient matrix A guarantee that the KP line-soliton solutions are regular. But no such clear prescription to obtain regular solutions is known for the Grammian case. This issue can be resolved by establishing an equivalence between the Wronskian and the Grammian forms for the KP τ -function. This will be done below. More specifically, it will be shown by explicit construction that there is a one-to-one correspondence between the formulas (7) and (11).

2.3. Equivalence of the Wronskian and Grammian forms of the τ -function

It is convenient to first start with the Wronskian form and derive the τ -function formula (11) from that. It follows from either (7) or (4) that the τ -function can be written as: $\tau = \det(AEK)$, where $E = \text{diag}(e^{\theta_1}, e^{\theta_2}, \dots, e^{\theta_M})$ is an $M \times M$ diagonal matrix and $K := (K_{ij}) = (k_i^{j-1})$ is an $M \times N$ Vandermonde matrix. In the following, the $N \times M$ coefficient matrix A will be chosen in RREF such that it can be represented as $A = (I, J)P$ with I and J denoting respectively, the $N \times N$ and $N \times (M - N)$ sub-matrices of the pivot and non-pivot columns of A , and P being the $M \times M$ permutation matrix which shuffles those columns to form A . For example, the coefficient matrix for the O-type 2-soliton τ -function discussed above can be represented as

$$A = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix} P_{23},$$

where P_{23} is the 4×4 matrix permuting the second and the third columns. Using the above form of A , the determinant for the τ -function becomes

$$\tau = |AEK| = |(I, J)(PEP^{-1})PK| = \left| (I, J) \begin{pmatrix} E_1 & 0 \\ 0 & E_2 \end{pmatrix} \begin{pmatrix} K_1 \\ K_2 \end{pmatrix} \right| = |IE_1K_1 + JE_2K_2|,$$

where E_1 and E_2 are respectively, $N \times N$ and $(M - N) \times (M - N)$ block diagonal matrices whose elements are permutations of the set $\{e^{\theta_1}, e^{\theta_2}, \dots, e^{\theta_M}\}$. Similarly, K_1 and K_2 are respectively, $N \times N$ and $(M - N) \times N$ matrices obtained by permuting the rows of the Vandermonde matrix K by P . It should be clear from above that in effect, the $M \times M$ matrix P induces a permutation π of the ordered set $\{k_1, k_2, \dots, k_M\}$ which can be expressed as

$$\pi(\{k_1, k_2, \dots, k_M\}) = \{q_1, q_2, \dots, q_N, p_1, p_2, \dots, p_{M-N}\}, \quad (12)$$

after renaming the elements of the permuted set. Accordingly, the matrices E_1, E_2, K_1, K_2 are redefined as

$$\begin{aligned} E_1 &= \text{diag}(e^{\phi(q_1)}, e^{\phi(q_2)}, \dots, e^{\phi(q_N)}), & E_2 &= \text{diag}(e^{\phi(p_1)}, e^{\phi(p_2)}, \dots, e^{\phi(p_{M-N})}), \\ K_1 &:= (K_{1ij})_{i,j=1}^N = (q_i^{j-1}), & K_2 &:= (K_{2ij}) = (p_i^{j-1}), \end{aligned}$$

for $i = 1, \dots, M - N$, $j = 1, \dots, N$, and where $\phi(k)$ is defined below (9). Since the parameters $q_1, \dots, q_N$ are distinct, the Vandermonde matrix K_1 is invertible. Then the above determinant expression for τ can be further manipulated as

$$\tau = |E_1 K_1| \hat{\tau}, \quad \hat{\tau} = |I + JE_2 K_2 K_1^{-1} E_1^{-1}|,$$

where τ and $\hat{\tau}$ differ by an exponential factor linear in x, y, t , hence both generate the same KP solution via (3). To show that $\hat{\tau}$ is indeed the Grammian form of the τ -function,

one employs the following matrix identity which is derived in the Appendix,

$$K_2 K_1^{-1} = D_2 \chi D_1^{-1}, \quad (13)$$

where the matrices D_1 , D_2 and the Cauchy matrix χ are defined as follows:

$$D_1 = \text{diag} \left(\prod_{j=1, j \neq i}^N (q_i - q_j) \right)_{i=1}^N, \quad D_2 = \text{diag} \left(\prod_{j=1}^N (p_i - q_j) \right)_{i=1}^{M-N},$$

$$\chi = (\chi)_{ij} = \frac{1}{p_i - q_j}, \quad i = 1, \dots, M - N, \quad j = 1, \dots, N.$$

Substitution of (13) into the expression for $\hat{\tau}$ above, yields

$$\hat{\tau} = |I + J D_2 E_2 \chi E_1^{-1} D_1^{-1}| = |I + (D_1^{-1} J D_2)(E_2 \chi E_1^{-1})|.$$

Finally, by setting $r = M - N$ and the $N \times M - N$ matrix $C = D_1^{-1} J D_2$, the Grammian form of the τ -function in (11) is readily recovered from the expression of $\hat{\tau}$ above.

It is relatively straightforward to reverse the steps described above to obtain the Wronskian data, i.e., the coefficient matrix A and the ordered set of parameters $k_1 < k_2 < \dots < k_M$ from the Grammian data, which consists of the $N \times M - N$ matrix C and the *unordered* set of parameters $\{q_1, \dots, q_N, p_1, \dots, p_{M-N}\}$. The key step is to recover the permutation π in (12) by simply *ordering* the set of Grammian parameters, i.e.,

$$\pi^{-1}(\{q_1, \dots, q_N, p_1, \dots, p_{M-N}\}) = \{k_1, k_2, \dots, k_M\}.$$

This in turn, provides the $M \times M$ permutation matrix P from π , and the matrix A is then constructed by the formula $A = (I, J)P$ with $J = D_1 C D_2^{-1}$, thus establishing the Grammian-Wronskian equivalence for the τ -function for the KP line-soliton solutions. Such equivalence was also derived in Ref. [25] for special N -soliton solutions using similar techniques.

An alternative approach to obtain the Wronskian form from the Grammian in (8) that is applicable when at least one of the functions f_i or g_j is an exponential, was discussed in [26]. In this case, one can for instance, choose the functions $\{f_i\}_{i=1}^N$ as N linearly independent solutions of (5), and $\{g_j := e^{-\theta_j}\}_{j=1}^N$ with $\theta_j = k_j x + k_j^2 y + k_j^3 t$ and $k_j > 0$. Using these forms of f_i and g_j in (8), and choosing Γ to be the path from $(-\infty, y)$ to (x, y) parallel to the x -axis, the integral for the matrix element Ω_{ij} can be expressed as an infinite series in inverse powers of the parameters k_j by repeated integration by parts, provided that $f_i^{(n)} g_j \rightarrow 0$, $n \geq 0$ as $x \rightarrow -\infty$. Then by taking the limit $k_j \rightarrow \infty$, $j = 1, \dots, N$, one can recover the Wronskian form (4) from the determinant $\det(\Omega)$.

An immediate consequence of the above equivalence result is that it is now possible to formulate the regularity condition for the KP line-solitons in a precise fashion from the Grammian form (11) of the τ -function. Given the matrices C and P , a necessary and sufficient condition that the KP solution (3) is non-singular is given by the fact that $A = (I, J)P$ be a totally non-negative matrix. Of course, this is the same condition on the coefficient matrix A in the Wronskian form. If in addition, A satisfies the following *irreducibility* conditions:

- (i) each column of A contains at least one nonzero element,
- (ii) each row of A in RREF contains at least one nonzero element other than the pivot (first non-zero entry),

then the number of line solitons as $y \rightarrow \pm\infty$ in the solution generated by the τ -function in (7) is determined by the size of the matrix A . Namely, one has $N_- = M - N$ and

$N_+ = N$, where $N_{\pm}$ denote the number of line solitons as $y \rightarrow \pm\infty$ [20, 21, 23]. Further analysis of the totally non-negative matrix A leads to the precise identification of the line solitons and a comprehensive classification scheme for all line-soliton solutions of the KP equation. The latter problem is related to the classification of the non-negative cells $\text{Gr}^+(N, M)$ of the Grassmannian $\text{Gr}(N, M)$ in terms of certain types of permutations called *derangements* [21].

It is worth noting that the equivalence between the Grammian and Wronskian forms of the line-soliton τ -function is not unique to the KP equation but applies also to other equations whose τ -functions are represented in both forms. Examples include several $2 + 1$ -dimensional differential-difference equations such as the 2d-Toda, 2d-Volterra and the differential-difference KP equation as well as difference-difference equations such as the fully discrete 2d-Toda and the discrete KP (Hirota-Miwa) equation. These will be discussed in future works.

3. O- and P-type 2-solitons

In this section the soliton interactions associated with certain types of 2-soliton solutions of the KP equation will be investigated. These solutions are characterized by a pair of line solitons as $|y| \rightarrow \infty$. Thus, $N_- = N_+ = 2$, which implies that $N = 2$ and $M = 4$. The Wronskian form of the τ -function for these solutions is obtained from (7) in terms of the ordered set $k_1 < k_2 < k_3 < k_4$ of distinct real parameters and a 2×4 totally non-negative matrix A as,

$$\tau(x, y, t) = \sum_{1 \leq r < s \leq 4} (k_s - k_r) A(r, s) e^{\theta_r + \theta_s}, \quad (14)$$

where $A(r, s)$ denotes the 2×2 minors of the matrix A . Taking into account the irreducibility property mentioned in Section 2, there are exactly *seven* distinct types of totally non-negative 2×4 matrices, which lead to seven different types of 2-soliton solutions of the KP equation [21, 23]. In contrast, its (1+1)-dimensional version namely, the KdV equation has only one kind of 2-soliton solution. This is indicative of a much richer solution space for the (2+1)-dimensional integrable equations than the (1+1)-dimensional ones.

The particular 2-soliton solutions whose nonlinear interaction properties are studied here, are the O-type 2-soliton which was mentioned in section 2, and another one called the P-type (“P” for *physical*), which fits better with the physical description of oblique interactions of shallow water waves for which the KP equation is a good approximation. In both of these cases, a pair of line solitons interact to form a X-shape in the xy -plane but the details of the interactions are entirely different. The corresponding coefficient matrices in RREF are given by

$$A_O = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}, \quad A_P = \begin{pmatrix} 1 & 0 & 0 & -1 \\ 0 & 1 & 1 & 0 \end{pmatrix},$$

which, from (14), lead to the τ -functions

$$\begin{aligned} \tau_O &= (k_3 - k_1)e^{\theta_1 + \theta_3} + (k_3 - k_2)e^{\theta_2 + \theta_3} + (k_4 - k_1)e^{\theta_1 + \theta_4} + (k_4 - k_2)e^{\theta_2 + \theta_4}, \\ \tau_P &= (k_2 - k_1)e^{\theta_1 + \theta_2} + (k_3 - k_1)e^{\theta_1 + \theta_3} + (k_4 - k_2)e^{\theta_2 + \theta_4} + (k_4 - k_3)e^{\theta_3 + \theta_4}. \end{aligned}$$

Applying the principle of dominant balance mentioned in the examples given in Section 2, it is possible to identify the line solitons in the two cases. For the O-type the line solitons




Figure 2. An O-type 2-soliton. The interaction peak corresponds to an absolute maximum.

are $[1, 2]$ and $[3, 4]$ as $y \rightarrow \pm\infty$, whereas for the P-type, these are $[1, 4]$ and $[2, 3]$. Recall that locally, a line soliton $[i, j]$ is given by (2) with parameters k_i and k_j .

For both O- and P-type solitons, the exact solution can be computed as

$$u = 2(\ln \tau)_{xx} = \frac{\alpha \operatorname{sech}^2 \phi_1 + \beta \operatorname{sech}^2 \phi_2 + \gamma \operatorname{sech}^2 \phi_1 \operatorname{sech}^2 \phi_2}{1 + C \tanh \phi_1 \tanh \phi_2},$$

with $\alpha = A_1(1 - C^2)$, $\beta = A_2(1 - C^2)$, and $\gamma = (A_1 + A_2)C^2 + 2C\sqrt{A_1 A_2}$. The expressions for the soliton amplitudes A_1, A_2 , soliton phases ϕ_1, ϕ_2 , and the constant C are different for the O- and P-type 2-solitons. These are given below.

3.1. O-type interaction

For the O-type, the amplitudes of the line solitons $[1, 2]$ and $[3, 4]$ are

$$A_1 = \alpha_{12} = \frac{1}{2}(k_2 - k_1)^2, \quad A_2 = \alpha_{34} = \frac{1}{2}(k_4 - k_3)^2,$$

the soliton phases are expressed in terms of $\theta_j = k_j x + k_j^2 y + k_j^3 t + \theta_{0j}$, $j = 1, \dots, 4$, as

$$\phi_1 = \frac{1}{2}(\theta_2 - \theta_1) + \frac{1}{4} \ln \left[\frac{(k_4 - k_2)(k_3 - k_2)}{(k_4 - k_1)(k_3 - k_1)} \right], \quad \phi_2 = \frac{1}{2}(\theta_4 - \theta_3) + \frac{1}{4} \ln \left[\frac{(k_4 - k_2)(k_4 - k_1)}{(k_3 - k_2)(k_3 - k_1)} \right],$$

and the constant C is given in terms of the parameters $k_1, \dots, k_4$ by

$$C = \frac{1 - \sqrt{\Delta_O}}{1 + \sqrt{\Delta_O}}, \quad \Delta_O = \frac{(k_3 - k_2)(k_4 - k_1)}{(k_3 - k_1)(k_4 - k_2)} = 1 - \frac{(k_2 - k_1)(k_4 - k_3)}{(k_3 - k_1)(k_4 - k_2)} < 1,$$

such that $0 < C < 1$. An O-type soliton is illustrated in Figure 2. The function $u(x, y, t)$ has an absolute maximum at $\phi_1 = \phi_2 = 0$, and the maximum value at this interaction point is given by (see [23])

$$u_{\max} = A_1 + A_2 + 2C\sqrt{A_1 A_2}. \quad (15)$$

Since $C > 0$, the interaction peak $u_{\max}$ is always greater than the sum of the individual line soliton amplitudes. In fact, one can easily verify that

$$A_1 + A_2 < u_{\max} < (\sqrt{A_1} + \sqrt{A_2})^2.$$




Figure 3. (a) Symmetric choice of the k -parameters: $k_4 = -k_1, k_3 = k = -k_2$; (b) a O-type 2-soliton with $k_4 = 2 = -k_1, k_3 = 1 = -k_2$. Here, each of the $[1, 2]$ and $[3, 4]$ line soliton makes an angle ψ with the y -axis.

Furthermore, $u_{\max}$ depends non-linearly on the incidence angle between the two line solitons. In order to investigate this behavior, it is convenient to choose the parameters $k_1, \dots, k_4$ such that $k_1 = -k_4$ and $k_2 = -k_3 = -k$ (see Figure 3(a)). Then the two line solitons are placed symmetrically about the y -axis, and are of equal amplitude, i.e., $A_1 = A_2 = A$. Denoting the angle between the $[3, 4]$ -soliton and the y -axis by ψ as shown in Figure 3(b), it follows that $\tan \psi = (k_3 + k_4) = \sqrt{2A} + 2k$. Since $k > 0$, the angle ψ is always greater than the critical angle

$$\psi_c = \tan^{-1}(\sqrt{2A}),$$

which depends on the amplitude of the line solitons. Setting $k_1 = -k_4, k_2 = -k_3$ in the expression for C above, it can be deduced from (15) that

$$u_{\max} = \frac{4A}{1 + \sqrt{1 - s^2}}, \quad s = \frac{\tan \psi_c}{\tan \psi} < 1$$

Therefore, $u_{\max}$ is a decreasing function of the angle ψ , and as $\psi \rightarrow \psi_c$ (i.e., $k \rightarrow 0$), the peak interaction amplitude $u_{\max} \rightarrow 4A$. Note that in the limit $k = 0$, the parameters $k_2 = k_3$, which implies that the τ -function τ_O contains only *three* instead of four exponential terms. The resulting solution $u(x, y, t)$ of the KP equation is a Miles' Y-shape solution corresponding to the confluence of three line solitons interacting resonantly [11] (see also [23]). The dependence of $u_{\max}$ with the angle of interaction is shown in Figure 4 for soliton amplitudes $A = 1$ and $A = 2$. Note that in Figure 4(a), $u_{\max} \rightarrow 4$ as ψ approaches the critical angle $\psi_c \approx 54.7^\circ$, while in Figure 4(b), $u_{\max} \rightarrow 8$ as $\psi \rightarrow \psi_c \approx 63.4^\circ$.




Figure 4. Plots of $u_{\max}$ vs ψ for O-type 2-solitons. The amplitudes of the line solitons $[1, 2]$ and $[3, 4]$ are fixed and equal, i.e., $A_1 = A_2 = A$: (a) $A = 1, \psi_c \approx 54.7^\circ$; (b) $A = 2, \psi_c \approx 63.4^\circ$.




Figure 5. (a) A P-type 2-soliton. The interaction between the line solitons is characterized by a saddle point; (b) Contour plot of a P-type 2-soliton with $k_1 = -3, k_2 = -0.5, k_3 = 2, k_4 = 3$. The line soliton $[1, 4]$ is along the y -axis, the angle between the line solitons $[1, 4]$ and $[2, 3]$ is ψ .

3.2. P-Type interaction

For the P-type, the soliton amplitudes

$$A_1 = \alpha_{14} = \frac{1}{2}(k_4 - k_1)^2, \quad A_2 = \alpha_{23} = \frac{1}{2}(k_3 - k_2)^2,$$

are always unequal, and $A_1 > A_2$. The soliton phases are given by

$$\phi_1 = \frac{1}{2}(\theta_4 - \theta_1) + \frac{1}{4} \ln \left[\frac{(k_2 - k_1)(k_3 - k_1)}{(k_4 - k_2)(k_4 - k_3)} \right], \quad \phi_2 = \frac{1}{2}(\theta_3 - \theta_2) + \frac{1}{4} \ln \left[\frac{(k_4 - k_2)(k_2 - k_1)}{(k_4 - k_3)(k_3 - k_1)} \right],$$

and the constant C is negative since

$$C = -\frac{1 - \sqrt{\Delta_P}}{1 + \sqrt{\Delta_P}}, \quad \Delta_P = \frac{(k_2 - k_1)(k_4 - k_3)}{(k_3 - k_1)(k_4 - k_2)} < 1.$$

A P-type 2-soliton solution is illustrated in Figure 5. In contrast to the O-type, the P-type 2-soliton solution $u(x, y, t)$ corresponds to *saddle* at the interaction point $\phi_1 = \phi_2 = 0$. The value of the solution at the interaction point is

$$u_{\text{int}} = A_1 + A_2 + 2C\sqrt{A_1 A_2}. \quad (16)$$

Since $C < 0$ for the P-type solution, u_{int} is always less than the sum of the amplitudes of the line solitons $[1, 4]$ and $[2, 3]$. A more precise bound for u_{int} is obtained as follows [23]: From the above expression for C , one can calculate

$$-C = \frac{1 - \sqrt{\Delta_P}}{1 + \sqrt{\Delta_P}} = \frac{\sqrt{(1-x)(1-y)} - \sqrt{xy}}{\sqrt{(1-x)(1-y)} + \sqrt{xy}} = \frac{1 - (x+y)}{(\sqrt{(1-x)(1-y)} + \sqrt{xy})^2},$$

where $x = \frac{k_2 - k_1}{k_4 - k_1}$ and $y = \frac{k_4 - k_3}{k_4 - k_1}$. Since $0 < x, y < 1$, the following inequalities

$$\sqrt{xy} \leq \frac{1}{2}(x+y), \quad \sqrt{(1-x)(1-y)} \leq 1 - \frac{1}{2}(x+y)$$

hold, and from these it can be easily deduced that $\sqrt{(1-x)(1-y)} + \sqrt{xy} \leq 1$. The latter inequality leads to

$$\frac{1 - \sqrt{\Delta_P}}{1 + \sqrt{\Delta_P}} \geq 1 - (x+y) = \frac{k_3 - k_2}{k_4 - k_1} = \sqrt{\frac{A_2}{A_1}}.$$

Thus, $-1 \leq C \leq -\sqrt{A_2/A_1}$, and from (16) it follows that

$$(\sqrt{A_1} - \sqrt{A_2})^2 < u_{\text{int}} \leq A_1 - A_2.$$

Next we investigate the dependence of the amplitude u_{int} at the interaction point on the angle between the two line solitons $[1, 4]$ and $[2, 3]$ with fixed amplitudes A_1 and A_2 . For simplicity, we take the $[1, 4]$ -soliton along the y -axis by setting $k_1 + k_4 = 0$, and denote by ψ the angle between the solitons as in Figure 5(b). Then

$$\tan \psi = k_2 + k_3 = 2(k_2 - k_1) - (\sqrt{2A_1} - \sqrt{2A_2}),$$

after using $k_1 + k_4 = 0$. Similarly, one calculates $(k_2 - k_1) + (k_4 - k_3) = \sqrt{2A_1} - \sqrt{2A_2}$, which implies that $0 < k_2 - k_1 < \sqrt{2A_1} - \sqrt{2A_2}$. Hence for fixed amplitudes A_1 and A_2 , the angle ψ between the two line solitons satisfies

$$-\psi_c < \psi < \psi_c, \quad \psi_c = \tan^{-1}(\sqrt{2A_1} - \sqrt{2A_2}).$$

The limiting cases $\psi = \pm\psi_c$ correspond to $k_1 = k_2$ or $k_3 = k_4$. Either case leads to a degeneration of the P-type 2-soliton to a Y-shape solution similar to O-type 2-soliton situation. In terms of A_1 , A_2 and the angle ψ , the quantity Δ_P can be expressed as

$$\Delta_P = \frac{(k_2 - k_1)(k_4 - k_3)}{(k_3 - k_1)(k_4 - k_2)} = \frac{\tan^2 \psi_c - \tan^2 \psi}{(\sqrt{2A_1} + \sqrt{2A_2})^2 - \tan^2 \psi},$$

and then from (16) one obtains

$$u_{\text{int}} = \frac{1}{2} \left[\tan^2 \psi + \sqrt{(\tan^2 \psi_c - \tan^2 \psi)((\sqrt{2A_1} + \sqrt{2A_2})^2 - \tan^2 \psi)} \right].$$

It follows from the above expression that u_{int} is an even function of the angle ψ between the line solitons $[1, 4]$ and $[2, 3]$ for $|\psi| < \psi_c$. When $\psi = 0$, i.e., when the two line solitons are both parallel to the y -axis, u_{int} reaches its maximum value of $A_1 - A_2$ which is the difference between the amplitudes of the two line solitons. As $\psi \rightarrow \pm\psi_c$, the interaction amplitude approaches its lower bound, i.e., $u_{\text{int}} \rightarrow \frac{1}{2} \tan^2 \psi_c = (\sqrt{A_1} - \sqrt{A_2})^2$. Figure 6 below illustrates the plots of u_{int} as a function of the angle ψ for fixed values of soliton amplitudes A_1 and A_2 .




Figure 6. Plots of u_{int} vs interaction angle ψ for P-type 2-solitons with fixed amplitudes A_1, A_2 of the line solitons $[1, 4]$ and $[2, 3]$: (a) $A_1 = 2, A_2 = 1$ and $\psi_c \approx 30.4^\circ$. The maximum value of u_{int} is 1 at $\psi = 0$, and $u_{\text{int}} \rightarrow (\sqrt{A_1} - \sqrt{A_2})^2 \approx 0.17$ as $\psi \rightarrow \pm\psi_c$; (b) $A_1 = 4, A_2 = 1$ and $\psi_c \approx 54.7^\circ$. The maximum interaction amplitude $(u_{\text{int}})_{\text{max}} = 3$ at $\psi = 0$, and $u_{\text{int}} \rightarrow 1$ as $\psi \rightarrow \pm\psi_c$.

4. Conclusion

In this article we have considered a special class of non-singular solutions of the KP equation referred to as the line-solitons which decay exponentially as $x, y \rightarrow \pm\infty$ except along certain directions in the xy -plane. Such solutions exhibit a variety of time-dependent spatial patterns due to resonant soliton interactions (Y-shape) as well as non-resonant interactions (X-shape). The exact analytic form of these solutions can be derived from the associated τ -functions which are expressible either as Wronskians or as Gram determinants. It is remarkable that the KP equation possesses such a rich structure of line-soliton solutions generated from a simple form of the τ -function. It turns out that the solution manifold of the line-solitons is parametrized by a discrete set of real distinct parameters and the space of totally non-negative matrices. This characterization is clear from the Wronskian form of the KP τ -function but not so transparent from its Grammian form. It is perhaps due to this difficulty in imposing appropriate regularity conditions that only a handful of line-soliton solutions were explicitly known via the direct algebraic methods which used the Grammian form of the τ -function. This issue has been resolved in this article where a one-to-one correspondence between the two forms of the line-soliton τ -function has been established. Consequently, it is now possible to derive non-singular line-soliton solutions using the Grammian form of the τ -function as well.

Another problem discussed in this article is the nonlinear soliton interactions for certain types of 2-soliton solutions. In particular, the amplitude of the nonlinear interaction has been explicitly calculated from the exact analytic expression for each of the 2-soliton solutions considered in this paper. Moreover, the dependence of the interaction amplitude on the angle between the line solitons has been investigated by keeping the line soliton amplitudes fixed. One possible physical significance of such results lies in the study of oblique nonlinear interactions of weakly 2-dimensional solitary waves in shallow water. The physical mechanism generating large amplitude waves of extreme elevations from the interaction of two (or more) smaller amplitude solitary waves in shallow water constitutes an important open problem. It is believed that in appropriate parameter regimes the line-solitons of the KP equation can serve as a reasonably good test-bed for the description and analysis of nonlinear solitary wave dynamics. As a qualitative evidence, one may consider the simple example of the O-type 2-soliton interaction where the interaction peak amplitude may reach as high as *four* times the amplitude of the individual line solitons.

For both O- and P-type solutions, the range of this interaction angle is found to be limited by a critical value ψ_c which depends on the amplitudes of the line solitons. It is however important to note that the theory of KP line-solitons can still be applied to study wave interaction where the angle between the incident waves is *outside* of the prescribed range for the O- or the P-type soliton solutions. In such cases, the wave dynamics is governed by other types of KP 2-soliton solutions which have recently been uncovered [17, 20, 21]. The study of the interaction properties of these newly found solutions is a topic of future investigation.

Acknowledgments

We thank Yuji Kodama for useful discussions. The research of the first and second authors is partially supported by the NSF grant DMS-0807404.

References

- [1] B. B. Kadomtsev and V. I. Petviashvili, *On the stability of solitary waves in weakly dispersive media*, Sov. Phys. - Dokl. 15 (1970), pp. 539–541.
- [2] E. Infeld and G. Rowlands, *Nonlinear waves, solitons and chaos*, Cambridge University Press, Cambridge, 2000.
- [3] S. Novikov, S. V. Manakov, L. P. Pitaevskii and V. E. Zakharov, *Theory of Solitons: The Inverse Scattering Method*, Contemporary Soviet Mathematics, Consultants Bureau, New York and London, 1984.
- [4] M. J. Ablowitz and P. A. Clarkson, *Solitons, nonlinear evolution equations and inverse scattering*, Cambridge University Press, Cambridge, 1991.
- [5] L. A. Dickey, *Soliton equations and Hamiltonian systems*, Advanced Series in Mathematical Physics Vol. 12, World Scientific, Singapore, 1991.
- [6] R. Hirota, *The Direct Method in Soliton Theory*, Cambridge University Press, Cambridge, 2004.
- [7] M. Sato, *Soliton equations as dynamical systems on an infinite dimensional Grassmannian manifold*, RIMS Kokyuroku (Kyoto University) 439 (1981), pp. 30–46.
- [8] V. E. Zakharov and A. B. Shabat, *A scheme for integrating nonlinear equations of mathematical physics by the method of the inverse scattering problem*, Func. Anal. Appl. 8 (1974), pp. 226–235.
- [9] J. Satsuma, *N-Soliton solution of the two-dimensional Korteweg-de Vries equation*, J. Phys. Soc. Jpn. 40 (1976), pp. 286–290.
- [25] N. C. Freeman and J. J. C. Nimmo, *Soliton-solutions of the Korteweg-de Vries and Kadomtsev-Petviashvili equations: the Wronskian technique*, Phys. Lett. A 95 (1983), pp. 1–3.
- [11] J. W. Miles, *Resonantly interacting solitary waves*, J. Fluid Mech. 79 (1977), pp. 171–179.
- [12] A. C. Newell and L. G. Redekopp, *Breakdown of Zakharov-Shabat theory and soliton creation*, Phys. Rev. Lett. 38 (1977), pp. 377–380.
- [13] N. C. Freeman, *Soliton interactions in two-dimensions*, Adv. Appl. Mech. 20 (1980), pp. 1–37.
- [14] K. Ohkuma and M. Wadati, *The Kadomtsev-Petviashvili equation: the trace method and the soliton resonances*, J. Phys. Soc. Jpn. 52 (1983), pp. 749–760.
- [15] M. Boiti, F. Pempinelli, A. K. Pogrebkov and B. Prinari, *Towards an inverse scattering theory for non-decaying potentials of the heat equation*, Inverse Problems 17 (2001), pp. 937–957.
- [16] E. Medina, *An N soliton resonance for the KP equation: interaction with change of form and velocity*, Lett. Math. Phys. 62 (2002), pp. 91–99.
- [17] G. Biondini and Y. Kodama, *On a family of solutions of the Kadomtsev-Petviashvili equation which also satisfy the Toda lattice hierarchy*, J. Phys. A: Math. Gen. 36 (2003), pp. 10519–10536.
- [18] O. Pashaev and M. Francisco, *Degenerate four virtual soliton resonance for the KP-II* Theor. Math. Phys. 144 (2005), pp. 1022–1029.
- [19] Y. Kodama, *Young diagrams and N-soliton solutions of the KP equation*, J. Phys. A: Math. Gen. 37 (2004), pp. 11169–11190.
- [20] G. Biondini and S. Chakravarty, *Soliton solutions of the Kadomtsev-Petviashvili II equation*, J. Math. Phys. 47 (2006), 033514i (26 pp).
- [21] S. Chakravarty and Y. Kodama, *Classification of the line-solitons of KP-II*, J. Phys. A: Math. Theor. 41 (2008), 275209 (33 pp).
- [22] S. Chakravarty and Y. Kodama, *A generating function for the N-soliton solutions of the Kadomtsev-Petviashvili II equation*, Contemp. Math. 471 (2008), pp. 47–67.
- [23] S. Chakravarty and Y. Kodama, *Soliton solutions of the KP equation and application to shallow water waves*, Stud. Appl. Math. 123 (2009), pp. 83–151.
- [24] V. B. Matveev and M. A. Salle, *Darboux Transformations and Solitons* Springer-Verlag, Berlin 1991.
- [25] N. C. Freeman and J. J. C. Nimmo, *Soliton solutions of the Korteweg-de Vries and Kadomtsev-Petviashvili equations: the wronskian technique*, Proc. R. Soc. Lond. A389 (1983), pp. 319–329.
- [26] V. M. Galkin, D. E. Pelinovsky and Yu. A. Stepanyants, *The structure of the rational solutions to the Boussinesq equation*, Physica D 80 (1995), pp. 246–255.

Appendix A.

Here we provide an elementary derivation of the matrix identity (13) involving the Vandermonde matrices K_1 and K_2 . Recall that K_1 is an $N \times N$ matrix with elements $(K_1)_{ij} = q_i^{j-1}$, and K_2 is an $N \times (M - N)$ matrix with elements $(K_2)_{ij} = p_i^{j-1}$ where $\{q_1, q_2, \dots, q_N, p_1, p_2, \dots, p_{M-N}\}$ is a set of distinct elements.

Consider N monic polynomials $P_1(x), P_2(x), \dots, P_N(x)$ such that each polynomial is of degree $N - 1$ and has *distinct* real roots which consist of $N - 1$ elements from the set $\{q_1, q_2, \dots, q_N\}$. These polynomials can be expressed as

$$P_j(x) := \prod_{k=1, k \neq j}^N (x - q_k) = \sum_{n=1}^N P_{nj} x^{n-1}, \quad j = 1, 2, \dots, N, \quad (\text{A1})$$

where P_{nj} is the coefficient of x^{n-1} in the polynomial $P_j(x)$, and $P_{Nj} = 1$. An immediate

consequence of (A1) is that the polynomials satisfy

$$P_j(q_i) = \sum_{n=1}^N P_{nj} q_i^{n-1} = (K_1 P)_{ij} = (D_1)_{ii} \delta_{ij}, \quad i, k = 1, 2, \dots, N, \quad (\text{A2})$$

where $P = (P_{nj})_{n,j=1}^N$ is the matrix of polynomial coefficients, D_1 is the diagonal matrix defined below (13), and δ_{ij} is the Kröner symbol. Equation (A2) gives the matrix equation $K_1 P = D_1$ which implies that

$$P = K_1^{-1} D_1 \quad (\text{A3})$$

since K_1 is invertible. Next, evaluating the polynomials in (A1) at $x = p_i$, yields

$$P_j(p_i) = \sum_{n=1}^N P_{nj} p_i^{n-1} = \prod_{k=1, k \neq j}^N (p_i - q_k) = \frac{\prod_{n=1}^N (p_i - q_n)}{p_i - q_j},$$

for $i = 1, \dots, M - N$, $j = 1, \dots, N$. These can be represented by the matrix equation

$$K_2 P = D_2 \chi, \quad (\text{A4})$$

where the diagonal matrix D_2 and the Cauchy matrix χ are also defined below (13). Combining (A3) and (A4) gives the desired identity (13).